

EpiData Analysis <http://www.epidata.dk> V1.0 implemented from build 149
Observation numbers used for percentiles with small N (N<20)
 e.g. for the 75% percentile of 12 observations the value of variable
 in question will be taken from observation 10

N	Percentil											
	0,010	0,025	0,050	0,100	0,250	0,500	0,750	0,900	0,950	0,975	0,990	
1	1	1	1	1	1	1	1	1	1	1	1	1
2	1	1	1	1	1	2	2	2	2	2	2	2
3	1	1	1	1	1	2	3	3	3	3	3	3
4	1	1	1	1	1	3	4	4	4	4	4	4
5	1	1	1	1	2	3	5	5	5	5	5	5
6	1	1	1	1	2	4	5	6	6	6	6	6
7	1	1	1	1	2	4	6	7	7	7	7	7
8	1	1	1	1	2	5	7	8	8	8	8	8
9	1	1	1	1	3	5	8	9	9	9	9	9
10	1	1	1	1	3	6	8	10	10	10	10	10
11	1	1	1	1	3	6	9	11	11	11	11	11
12	1	1	1	1	3	7	10	12	12	12	12	12
13	1	1	1	1	4	7	11	13	13	13	13	13
14	1	1	1	2	4	8	11	14	14	14	14	14
15	1	1	1	2	4	8	12	14	15	15	15	15
16	1	1	1	2	4	9	13	15	16	16	16	16
17	1	1	1	2	5	9	14	16	17	17	17	17
18	1	1	1	2	5	10	14	17	18	18	18	18
19	1	1	1	2	5	10	15	18	19	19	19	19

This table used for all percentiles in commands where relevant